

Quality of life in painful diabetic neuropathy patients following treatment with spinal cord stimulation

Duarte, Rui; de Vos, Cecile; Andronis, Lazaros; Lenders , Mathieu

DOI:
[10.1111/ner.12333](https://doi.org/10.1111/ner.12333)

License:
None: All rights reserved

Document Version
Peer reviewed version

Citation for published version (Harvard):

Duarte, R, de Vos, C, Andronis, L & Lenders , M 2015, 'Quality of life in painful diabetic neuropathy patients following treatment with spinal cord stimulation', International Neuromodulation Society (INS) 12th World Congress, Montreal, Quebec, Canada, 6/06/15 - 11/06/15 pp. e333. <https://doi.org/10.1111/ner.12333>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:
Eligibility for repository: Checked on 07/12/2015

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Quality of life in painful diabetic neuropathy patients following treatment with spinal cord stimulation

Rui V. Duarte,¹ Cecile C. de Vos,^{2,3} Lazaros Andronis,⁴ Mathieu W.P.M. Lenders ³

¹ School of Health and Population Sciences, University of Birmingham, Birmingham, UK

² Department of Clinical Neurophysiology, University of Twente, Enschede, The Netherlands

³ Department of Neurosurgery, Medisch Spectrum Twente, Enschede, The Netherlands

⁴ Health Economics Unit, University of Birmingham, Birmingham, UK

Introduction

Diabetes mellitus is a chronic condition with an increasing prevalence estimated to be 4.4% worldwide in 2030, the equivalent to 366 million people. Around one-third of diabetic patients are expected to develop painful diabetic neuropathy (PDN), which is the most disabling and costly complication of diabetes.

Spinal cord stimulation (SCS) is a recognised neuromodulation technique for the management of neuropathic pains such as complex regional pain syndrome ¹ and failed back surgery syndrome. ² Recently it has been reported that SCS results in significantly reduced pain and improved quality of life in patients with refractory painful diabetic neuropathy.³ The quality of life (QoL) analysis was based on the EQ-5D visual analogue scale scores. We have now evaluated QoL based on index values of the EQ-5D for the countries where this trial was conducted.

Methods

The methods used, patient demographics and results from the randomised controlled trial are reported elsewhere.³ Quality of life was assessed using the EQ-5D questionnaire. Health-related quality of life was derived from participants' responses to the widely used EuroQoL EQ-5D-3L questionnaire. Responses were converted into single (utility) indices using the Dutch tariff. Paired samples t-test were conducted to evaluate within group QoL changes. An ANCOVA analysis was undertaken to compare QoL between the groups while adjusting for the baseline score. Changes in sub-categories of the EQ-5D-3L were evaluated through the Mann-Whitney test for between-group analyses and the Wilcoxon signed-rank test for within-group analyses. Statistical analyses were conducted using the statistical software STATA (Release 13; College Station, TX: StataCorp LP).

Results

No statistically significant improvements were observed for the CMP group between baseline and six-month follow-up for the VASPI, EQ-5D utility or EQ-5D VAS scores (Table 1). Statistically significant improvements were observed for all outcome measures for the patients in the SCS group between baseline and six-month follow-up. Patients randomised to SCS experience greater pain relief and grater improvement in QoL as measured by the EQ-5D utility scores and EQ-5D VAS than those patients randomised to CMP. The interaction between groups and QoL baseline score was not significant $F(1,50)=2.391$, $p=0.128$. The ANCOVA comparing QoL between groups while adjusting for the baseline score was significant $F(1,50) = 14.274$, $p<0.001$. On the EQ-5D sub-categories, at six-months the patients randomised to SCS reported significant improvements in four out of five dimensions: mobility, usual activities, pain/discomfort and anxiety/depression when compared to baseline (Figure 1). Statistically significant differences were observed between groups for the pain/discomfort EQ-5D dimension.

Table 1. Pain and EQ-5D index scores the SCS and CMP treatment groups				
	SCS (n = 40) (mean, SD)		CMP (n = 20) (mean, SD)	
	Baseline	Six-months	Baseline	Six-months
Pain VAS, mean (SD)	74.94 (12.15)	29.22 (27.49) **	66.94 (18.48)	66.44 (21.86) ^^^
EQ-5D, mean (SD)	0.25 (0.27)	0.65 (0.28) **	0.44 (0.31) ^^	0.44 (0.33) ^
EQ-5D VAS, mean (SD)	49.17 (19.07)	60.89 (23.38) *	47.12 (16.11)	40.53 (20.35)^^^
CMP, conventional medical practice; SCS, spinal cord stimulation; VAS, visual analogue scale				
*p < .05, **p < .0001 (within a group), ^p < .05, ^^p < .01, ^^^p < .0001 (between groups)				




Figure 1. Comparison of proportion of patients reporting some problems or extreme problems in the EQ-5D subcategories

Conclusion

SCS has the potential to be an alternative for the management of refractory PDN. In addition to the significant reduction in pain observed in the de Vos et al RCT, SCS also appears to lead to significant improvements in QoL in this population. An economic evaluation is now in progress to evaluate the cost-effectiveness of SCS for the management of refractory PDN.

References

1. Kemler MA, Barendse GA, van Kleef M, de Vet HC, et al. N Engl J Med 2000; 343: 618-24.
2. Kumar K, Taylor RS, Jacques L, Eldabe S, et al. Pain 2007; 132: 179-88.
3. de Vos CC, Meier K, Zaalberg PB, Nijhuis HJ, et al. Pain 2014; 155: 2426-31.